

Symmetry Breaking in Induced-Charge Electrophoresis

Ehud Yariv*

Faculty of Mechanical Engineering, Technion — Israel Institute of Technology, Haifa 32000, Isreal

The electrophoretic motion of a conducting particle, driven by an induced-charge mechanism, is analyzed. The dependence of the motion upon particle shape is embodied in four tensorial coefficients that relate the particle velocities to the externally-applied electric field. Several families of particle shapes, whose members are unaffected by the field, are identified via use of symmetry arguments. Other particles translate and/or rotate in response to the imposed field, even if their net electric charge vanishes. The coefficients are represented as surface integrals of the electric potential over the particle boundary, thereby eliminating the need to solve the flow field.

Traditionally, the term electrophoresis applies to the motion of colloidal particles through an electrolyte solution. This motion results from the interaction between an applied electric field, say $\mathbf{E}_\infty$, and the net ionic charge accommodated in the Debye cloud surrounding the particle boundary (this charge being equal in magnitude and opposite in sign to that adsorbed at the particle surface). In many situations, the Debye-layer thickness is small compared with particle size. The electrokinetic flow is then described by an approximated equation set, obtained via use of singular perturbations [1]. In that asymptotic description, the Debye screening length assumes zero size. Thus, the electric potential φ and velocity field $\mathbf{v}$ in the electrically-neutral fluid domain are respectively governed by Laplace's and Stokes' equations, whereas the presence of the layer is reflected by the boundary conditions that apply at the particle boundary $\mathcal{S}$. These consist of the no-flux condition, $\hat{\mathbf{n}}(\mathbf{x}_s) \cdot \mathbf{E}(\mathbf{x}_s) = 0$, governing the normal component of the electric field, and the slip condition, $\mathbf{v}_s(\mathbf{x}_s) = -(\varepsilon/\mu) \zeta(\mathbf{x}_s) \mathbf{E}(\mathbf{x}_s)$, governing the relative fluid-particle motion. These conditions respectively reflect the particle impermeability to ionic current and the net action of the electric field on the charged Debye layer. Here, $\hat{\mathbf{n}}$ is an outward-pointing unit vector normal to $\mathcal{S}$, $\mathbf{x}_s$ is a position vector located on $\mathcal{S}$, ε and μ respectively denote the permittivity and viscosity of the solution, the "zeta potential" ζ constitutes the local value of the potential jump across the Debye layer, and $\mathbf{E} = -\nabla\varphi$ is the electric field.

The preceding description is adequate for dielectric media, capable of carrying bound electric charge. Electrokinetic flow can also be generated about a conducting body; such flow, however, is driven by a polarization mechanism, made possible by the charge mobility. As in the dielectric case, the electric charge is also situated on the surface; this, however, is not the result of any physicochemical bonding, but rather a simple consequence of Gauss law [2]. Moreover, the surface charge in a conductor distributes over its surface so as to guarantee zero electric field in its interior. The resulting dependence of the zeta potential upon $\mathbf{E}_\infty$ gives rise to a nonlinear slip mechanism, which is responsible for a rich variety of physical phenomena. The most notable feature

in this induced electrokinetic effect is the generation of electrokinetic flow about an initially-uncharged particle, this flow having no counterpart in conventional linear electrophoresis.

Flows driven by this induced mechanism, addressing both DC and AC external fields, have been discussed mainly in the Ukrainian literature (see [6]). A detailed description of induced-charge electrokinetics appeared in a recent paper by Squires and Bazant [3]. The authors, discussing a broader context of flows about polarizable media, coined the terms "induced-charge electro-osmosis" (ICEO) and "induced-charge electrophoresis" (ICEP) to respectively describe the associated fluid and particle motion. They also suggested [4] that the combination of nonlinear slip with symmetry breaking can lead to useful configurations which can perform a variety of microfluidic operations (such as pumping and mixing). Their suggestions focus upon electro-osmotic flows relative to fixed conducting elements (which may be held at a constant electric potential), rather than electrophoretic motion of freely-suspended particles, which is addressed herein. Nevertheless, electrophoretic analyses tend to provides insight to the inverse problem of electro-osmosis.

The simplest ICEO situation is that of an initially uncharged conducting particle which is positioned in a uniform and constant applied field $\mathbf{E}_\infty$. During a short time interval, a Faradaic current charges the Debye layer in proximity to the particle surface, with an equal and opposite charge being set up on the surface itself. The induced potential difference across the layer, namely the zeta potential, is proportional to $\mathbf{E}_\infty$. Accordingly, the electrokinetic slip that occurs at $\mathcal{S}$ is quadratic in $\mathbf{E}_\infty$. Given the linearity of the flow problem, so are all the variables describing the resulting fluid (and perhaps particle) motion.

The prototypical ICEO problem, which was investigated in [5] as a model for the electrokinetic flow about mercury drops, entails a spherical particle. Owing to the fore-aft symmetry of this flow [6], no hydrodynamic force (or torque) is exerted on the particle. This dipolar symmetry is obviously absent in flows about more general particle shapes. In principle, any deviation from a spherical shape may lead to hydrodynamic force and torque,

with a consequent particle motion. Indeed, such motion has been experimentally observed [7].

The purpose of this paper is to elucidate the interesting properties of this nonlinear phenomenon. I consider here a relatively simple case, addressing the electrophoretic motion of a freely-suspended conducting particle in a uniform and constant externally-applied field. (The present results are easily generalized to the case of AC fields.) The analysis begins with the case of an initially-uncharged particle and is followed by discussion of non-zero net charge effects.

Consider an uncharged conducting particle of an arbitrary shape and of linear dimension a , which is positioned in an electrolyte solution in the presence of an externally-applied constant electric field, $\mathbf{E}_\infty = E_\infty \hat{\mathbf{E}}$ ($\hat{\mathbf{E}}$ being a unit vector in the direction of $\mathbf{E}_\infty$). When describing the governing field equations, the position vector is conveniently measured from an (arbitrary) point O attached to the particle. It is also convenient to describe the particle motion by the instantaneous velocity of point O , $\mathbf{U}_O$, together with its angular velocity $\mathbf{\Omega}$.

For conducting particles which are impermeable to the current carrying ions (as is usually the case) the no-flux condition, $\hat{\mathbf{n}} \cdot \nabla \varphi = 0$, still holds. Thus, the electrostatic problem governing φ consists of this condition, together with Laplace's equation in the fluid domain, $\nabla^2 \varphi = 0$, and the far field condition, $\nabla \varphi \rightarrow -\mathbf{E}_\infty$. This problem (which also describes the field about a comparable dielectric particle) is linear and homogeneous in φ and $\mathbf{E}_\infty$, from which it follows that φ must be a linear function of $\mathbf{E}_\infty$. The attendant flow field is governed by the Stokes equations, the slip condition at $\mathcal{S}$, $\mathbf{v} = \mathbf{U}_O + \mathbf{\Omega} \times \mathbf{x} + (\varepsilon \zeta / \mu) \nabla \varphi$, and the attenuation condition at large distances from $\mathcal{S}$, $\mathbf{v} \rightarrow \mathbf{0}$. For a freely suspended particle, the requirement of vanishing hydrodynamic force and torque completely determines the flow field, as well as the values of $\mathbf{U}_O$ and $\mathbf{\Omega}$.

As in common in the literature (see [3]), it is assumed that the zeta potential is sufficiently low to be proportional to the local surface charge density q ,

$$\zeta = \frac{\lambda}{\varepsilon} q. \quad (1)$$

Here, λ is the Debye-layer thickness of the Debye layer. With no loss of generality, the uniform value of the electric potential in the particle interior is chosen as zero. Accordingly, the zeta potential is related to the potential value just outside the layer, $\zeta = -\varphi|_{\mathcal{S}}$. It is the nonlinear slip term $\varphi \nabla \varphi$ which is responsible for the peculiar characteristics of ICEP. Since the slip is the driver of the flow, this term results in a quadratic dependence of $\mathbf{v}$ upon $\mathbf{E}_\infty$. Owing to the linearity of the flow problem, the particle translational and rotational velocities must




FIG. 1: An axisymmetric particle possessing fore-aft symmetry: in the absence of net charge, the particle remains stationary; in the presence of an initial charge, the particle translates but does not rotate.

possess the following structure:

$$\mathbf{U}_O = \frac{\varepsilon a}{\mu} \mathbf{C}_O : \mathbf{E}_\infty \mathbf{E}_\infty, \quad (2a)$$

$$\mathbf{\Omega} = \frac{\varepsilon}{\mu} \mathbf{D} : \mathbf{E}_\infty \mathbf{E}_\infty. \quad (2b)$$

Here, $\mathbf{C}_O$ is a dimensionless third-order tensor (which depends upon the position of O) and $\mathbf{D}$ is a dimensionless third-order pseudo-tensor. These objects are intrinsic geometric properties of the particle, independent of its size and orientation relative to the imposed field. In a sense, they constitute a lumped description of the electrophoretic motion.

It is actually possible to demonstrate a variety of symmetry properties, expressed through the coefficients $\mathbf{C}_O$ and $\mathbf{D}$, for various families of particle shapes. These properties may be obtained without the need to explicitly solve the governing field equations. The simplest example involves isotropic particles (such as spheres). For that family, no vector (or tensor) can specify the particle geometry. Since no isotropic third-order tensor exists [8], it is impossible to construct a candidate for $\mathbf{C}_O$. Moreover, the only isotropic third-order pseudo-tensor is ϵ , the alternating triadic, which yields zero when contracted with the symmetric dyadic $\mathbf{E}_\infty \mathbf{E}_\infty$. Accordingly, isotropic particles neither translate nor rotate.

A larger family is that of all axisymmetric particles possessing fore-aft symmetries, such as spheroids and finite cylinders of circular cross section (see Fig. 1). These particles are characterized by a single vector $\mathbf{e}$ attached to their axis of symmetry. The only third-order tensors which may be constructed using $\mathbf{e}$ are $\mathbf{eee}$ and various permutations of $\mathbf{le}$ ($\mathbf{l}$ being the idem-factor). These tensors, however, are odd functions of $\mathbf{e}$, and do not satisfy the invariance to reversal of $\mathbf{e}$ required from a body possessing a fore-aft symmetry. Moreover, it is impossible to construct a third-order pseudo-tensor using the single vector $\mathbf{e}$. Accordingly, such particles behave like spheres: they neither translate nor rotate.

It appears that the generation of particle motion requires a stronger symmetry breaking. Consider the case of axisymmetric particles lacking fore-aft symmetry, such as cones and hemispheres. Again, these particles are

characterized by a single vector $\mathbf{e}$ attached to their axis of symmetry, whence the only candidate for $\mathbf{D}$ is again $\mathbf{e}$, which means that such particles do not rotate. Both $\mathbf{U}_O$ and $\mathbf{C}_O$ are therefore independent of O , and may be respectively denoted by $\mathbf{U}$ and $\mathbf{C}$. As with the previous family, the only candidates for $\mathbf{C}$ are $\mathbf{e}\mathbf{e}\mathbf{e}$ and various permutations of $\mathbf{le}$. For bodies lacking fore-aft symmetry, $\mathbf{C}$ is not necessarily invariant to reflection. Thus, such particles may translate (without rotation) under the action of an external field.

In principle, both $\mathbf{C}_O$ and $\mathbf{D}$ are obtainable from the detailed knowledge of the velocity field. In what follows, integral expressions for these tensors are derived in terms of the electric potential distribution. These expressions are calculated using the Lorentz reciprocal theorem [9], thus avoiding the formidable calculation of the flow problem. For simplicity of presentation, this procedure is demonstrated for non-rotating (e.g., axisymmetric) particles, for which the angular momentum balance is automatically satisfied (the generalization to arbitrary particles is trivial). In what follows, it is convenient to employ a dimensionless notation, wherein length variables are normalized with a , velocities with $\varepsilon a E_\infty^2 / \mu$, forces with $\varepsilon a^2 E_\infty^2$, stresses with εE_∞^2 , and the electric potential with $a E_\infty$.

Using this notation, the slip condition on $\mathcal{S}$ appears as $\tilde{\mathbf{v}} = \tilde{\mathbf{U}} - \tilde{\varphi} \tilde{\nabla} \tilde{\varphi}$ ($\tilde{\cdot}$ denotes here the dimensionless counterpart of the relevant physical quantity). The linear flow problem is decomposed into two separate parts. The first describes pure translation of the particle with a velocity $\tilde{\mathbf{U}}$ (in the absence of any slip); in the second the particle is held fixed, and the flow is generated by the imposed slip on $\mathcal{S}$, $\tilde{\mathbf{v}} = -\tilde{\varphi} \tilde{\nabla} \tilde{\varphi}$. In both problems the velocity field satisfies the Stokes equations and attenuates at large distances from the particle.

The force resulting from the first flow field is given by a Stokes drag expression, $\mathbf{F}_I = -\mathbf{K} \cdot \tilde{\mathbf{U}}$, where the positive-definite dimensionless translation tensor $\mathbf{K}$ is an intrinsic property of the particle geometry [9]. The force resulting from the second flow field is provided by the following quadrature:

$$\mathbf{F}_{II} = - \oint_{\mathcal{S}} d\tilde{A} \tilde{\mathbf{n}} \cdot \boldsymbol{\Sigma}^\dagger \cdot \tilde{\nabla} \tilde{\varphi} \quad (3)$$

which follows from a variant of the reciprocal theorem [10]. Here, $d\tilde{A}$ is a dimensionless area element (normalized with a^2) and the (dimensionless) “translational” stress triadic field $\boldsymbol{\Sigma}$ (defined in [9]) depends only upon the particle geometry and the position vector of the fluid point on $\mathcal{S}$: explicitly, Σ_{ijk} denotes the ij component of the stress tensor that would result from a pure translation of the particle with a unit velocity in the k direction. (The superscript $\dagger$ signifies right transposition.)

The requirement of a force-free particle, $\mathbf{F}_I + \mathbf{F}_{II} = \mathbf{0}$,

yields the following result:

$$\tilde{\mathbf{U}} = -\mathbf{K}^{-1} \cdot \oint_{\mathcal{S}} d\tilde{A} \tilde{\mathbf{n}} \cdot \boldsymbol{\Sigma}^\dagger \cdot \tilde{\varphi} \tilde{\nabla} \tilde{\varphi}. \quad (4)$$

Owing to the linearity of the electrostatic problem, it is clear that $\tilde{\varphi} = \boldsymbol{\Phi}(\tilde{\mathbf{x}}) \cdot \tilde{\mathbf{E}}$, where $\boldsymbol{\Phi}$, a dimensionless vector function of position, can only depend upon particle shape. Use of the definition (2a) furnishes the requisite representation,

$$\mathbf{C} = -\mathbf{K}^{-1} \cdot \oint_{\mathcal{S}} d\tilde{A} \tilde{\mathbf{n}} \cdot \boldsymbol{\Sigma}^\dagger \cdot \tilde{\nabla} \boldsymbol{\Phi} \boldsymbol{\Phi}. \quad (5)$$

It is emphasized that this result holds only for a particle which does not possess an initial charge.

Consider now a particle possessing an initial charge Q (or, equivalently, a net charge Q). In this situation, the zeta potential is not quadratic in $\mathbf{E}_\infty$ and the relations (2) are invalidated. Define an “effective” potential, $\psi \stackrel{\text{def}}{=} Q\lambda/A\varepsilon$, where $A = \oint_{\mathcal{S}} dA$ is the particle surface area. While it is customary to identify ψ with the zeta potential [cf. (1)], I here avoid any such *a priori* interpretation.

The flow about an initially-charged particle under the action of an applied field has been discussed in [3]. Given the linear-capacitor approximation (1), additivity in charge implies additivity in the zeta potential. Accordingly, the initially-charged situation was represented as a combination of two different flows: the first is associated with a quadratic slip condition, and the second is that involved in a linear electrophoretic motion corresponding to a uniform zeta potential $\zeta = \psi$. It is important to note, however, that the situation discussed in [3] — with a view towards microfluidic devices — is focused upon the specific case of a circular cylinder (a similar representation is also valid for the case of spheres). Is that interpretation valid for more general shapes?

The additional scale ψ modifies the translational and rotational relations (2), which, following dimensional arguments, are replaced by:

$$\mathbf{U}_O = \frac{\varepsilon a}{\mu} \mathbf{A}_O : \mathbf{E}_\infty \mathbf{E}_\infty + \frac{\varepsilon}{\mu} \psi \mathbf{M}_O \cdot \mathbf{E}_\infty + \frac{\varepsilon}{\mu a} \psi^2 \mathbf{P}_O \quad (6a)$$

$$\boldsymbol{\Omega} = \frac{\varepsilon}{\mu} \mathbf{B} : \mathbf{E}_\infty \mathbf{E}_\infty + \frac{\varepsilon}{\mu a} \psi \mathbf{N} \cdot \mathbf{E}_\infty + \frac{\varepsilon}{\mu a^2} \psi^2 \mathbf{R}. \quad (6b)$$

The dimensionless objects $\mathbf{A}_O$, $\mathbf{M}_O$, and $\mathbf{P}_O$ respectively denote a triadic, dyadic, and a vector, whereas the dimensionless objects $\mathbf{B}$, $\mathbf{N}$, and $\mathbf{R}$ respectively denote a pseudo-triadic, pseudo-dyadic, and a pseudo-vector. All six quantities are intrinsic geometric properties of the particle, independent of the applied field and the value of ψ .

In the case $\mathbf{E}_\infty = \mathbf{0}$ the translational and rotational velocities are respectively proportional to $\mathbf{P}_O$ and $\mathbf{R}$. Since the applied field is the driver of the motion, $\mathbf{P}_O$ and $\mathbf{R}$ must vanish in that case. These quantities, however, are independent of $\mathbf{E}_\infty$, and must therefore vanish identically. Similar logic can be applied via the case $\psi = 0$,

which yields relations similar to (2). Since $\mathbf{A}_O$ and $\mathbf{B}$ are independent of ψ , they must be respectively identical to the coefficients $\mathbf{C}_O$ and $\mathbf{D}$, which describe the motion of particles possessing zero net charge.

The role of $\mathbf{M}_O$ and $\mathbf{N}$ is elucidated via the asymptotic limit $aE_\infty/\psi \ll 1$, which corresponds to the quadratic effect (associated with the action of $\mathbf{E}_\infty$ on charge induced by $\mathbf{E}_\infty$) being negligible relative to the linear one (associated with the action of $\mathbf{E}_\infty$ on the initial charge distribution). In this case (6a)-(6b) degenerate to:

$$\mathbf{U}_O = \frac{\varepsilon}{\mu} \psi \mathbf{M}_O \cdot \mathbf{E}_\infty, \quad (7a)$$

$$\mathbf{\Omega} = \frac{\varepsilon}{\mu a} \psi \mathbf{N} \cdot \mathbf{E}_\infty. \quad (7b)$$

These are identical to the mobility relations governing conventional electrophoresis (with no induced charge effects), except for an important difference which will be identified shortly. Since $\mathbf{M}_O$ and $\mathbf{N}$ are independent of ψ and $\mathbf{E}_\infty$ (and, specifically, of the above-mentioned limit), they may be obtained using a linear electrophoretic analysis. It is therefore tempting to say (see [3]) that the mobility relations (7) are the same as those for a linear electrophoresis of particles having a zeta potential ψ , which would imply that $\mathbf{M}_O$ and $\mathbf{N}$ are governed by the Smoluchowski relation $\mathbf{U} = (\varepsilon\psi/\mu) \mathbf{E}_\infty$ and $\mathbf{\Omega} = \mathbf{0}$. That would yield $\mathbf{M} = \mathbf{I}$ and $\mathbf{N} = \mathbf{0}$.

It is important, however, to emphasize that the linear electrophoretic motion in the limit $aE_\infty/\psi \ll 1$ corresponds to a charge distribution in a *conducting* particle in the absence of any imposed field. Only for the case of a sphere (or a circular cylinder in two-dimensional flows) this distribution is uniform, corresponding to a uniform value (namely ψ) of the zeta potential. Thus, $\mathbf{M}_O$ and $\mathbf{N}$ would not be identical in general to those predicted by the Smoluchowski relations. (Indeed, linear electrophoresis of non-uniformly charged dielectric particles [11] is qualitatively different than that observed in the uniform case.) It appears that the traditional focus on spherical and cylindrical bodies leads to a loss of physical effects which characterize nonlinear slip. As such, conclusions drawn from the analyses of such specific shapes are rather restrictive and non-representative.

As in the case of particles possessing no net charge, for which the relations (2) apply, it is possible to deduce symmetry properties for the motion (namely the values of $\mathbf{M}_O$ and $\mathbf{N}$) of initially-charged particles. With axisymmetric particles, for example, $\mathbf{M}_O$ must be a linear combination of $\mathbf{I}$ and $\mathbf{ee}$, whereas $\mathbf{N}$ must be proportional to $\mathbf{e} \cdot \mathbf{e}$. In the low-field limit, $aE_\infty/\psi \ll 1$, such particles do not rotate, but in general may translate.

It is a simple matter to represent both $\mathbf{M}_O$ and $\mathbf{N}$ as quadratures of the electric field, without the need to solve the flow problem. As with the preceding analysis, which has led to the representation (5), the present derivation focuses upon non-rotating (e.g., axisymmetric) particles,

for which $\mathbf{N} = \mathbf{0}$ and $\mathbf{M}_O$ is independent of O (again, the generalization to the general case is trivial). In the limit $aE_\infty/\psi \ll 1$ the slip velocity is given by $(\varepsilon\zeta/\mu) \nabla\varphi$, with ζ being the zeta potential associated with the initial charge distribution (which is a function of $\mathbf{x}_s$). It is convenient to use the previously employed dimensionless notation, with which the present slip condition appears as $\tilde{\mathbf{v}} = \tilde{\mathbf{U}} + (\psi/aE_\infty) \tilde{\zeta} \tilde{\nabla}\tilde{\varphi}$ (with $\tilde{\zeta} = \zeta/\psi$). The concomitant flow field is decomposed into two parts, the first describing pure translation with a velocity $\tilde{\mathbf{U}}$, and the second describing a stationary particle on which the velocity slips with the prescribed value $(\psi/aE_\infty) \tilde{\zeta} \tilde{\nabla}\tilde{\varphi}$. A similar procedure to that leading to (4) then yields

$$\tilde{\mathbf{U}} = \frac{\psi}{aE_\infty} \mathbf{K}^{-1} \cdot \oint_S d\tilde{A} \tilde{\zeta} \tilde{\mathbf{n}} \cdot \mathbf{\Sigma}^\dagger \cdot \tilde{\nabla}\tilde{\varphi}. \quad (8)$$

Use of the linear representation, $\tilde{\varphi} = \mathbf{\Phi} \cdot \hat{\mathbf{E}}$, together with the definition (7a) furnishes the desired expression,

$$\mathbf{M} = \mathbf{K}^{-1} \cdot \oint_S d\tilde{A} \tilde{\zeta} \tilde{\mathbf{n}} \cdot \mathbf{\Sigma}^\dagger \cdot \tilde{\nabla}\mathbf{\Phi}. \quad (9)$$

It is re-emphasized that the zeta-potential distribution appearing in (9) corresponds to a surface charge arrangement on a *conducting* particle in a vacuum (in the absence of any applied field), whereas the electric field is identical to that about a *dielectric* particle in a vacuum.

For a sphere of radius a , the initial charge spreads uniformly, giving rise to a charge density Q/A . Thus, ψ is identical to the zeta potential, and $\tilde{\zeta} \equiv 1$. The potential distribution under the action of an applied external field $\mathbf{E}_\infty$ is given by a combination of a uniform field and a dipole, $\varphi = -(1 + a^3/2r^3) \mathbf{x} \cdot \mathbf{E}_\infty$, from which it follows that $\tilde{\nabla}\mathbf{\Phi}|_S = (3/2)(\hat{\mathbf{n}}\hat{\mathbf{n}} - \mathbf{I})$. Also, from the solution of the Stokes equations for a translating sphere [9], it is known that $\hat{\mathbf{n}} \cdot \mathbf{\Sigma} = -(3/2)\mathbf{I}$ and $\mathbf{K} = 6\pi\mathbf{I}$. Use of the integral relations $\oint_S d\tilde{A} = 4\pi$ and $\oint_S d\tilde{A} \tilde{\mathbf{n}}\hat{\mathbf{n}} = (4\pi/3)\mathbf{I}$ readily yields the Smoluchowski relation $\mathbf{M} = \mathbf{I}$.

I thank Martin Z. Bazant for turning my attention to ICEO, and for his valuable help in obtaining the Russian references. I thank Todd M. Squires for some useful discussions.

* Electronic address: yarive@technion.ac.il

- [1] R. W. O'Brien, J. Colloid Interface Sci. **92**, 204 (1983).
- [2] This would be true even in unsteady situations, the reason being the large ratio between typical conductivities of metallic conductors (say 6×10^7 S/m for copper) to those characterizing ionic solutions (where the conductivity, which depends upon the ionic concentration, usually does not exceed 10^{-6} S/m).
- [3] T. M. Squires and M. Z. Bazant, J. Fluid Mech. **509**, 217 (2004).
- [4] M. Z. Bazant and T. M. Squires, Phys. Rev. Lett. **92** (2004).

- [5] V. G. Levich, *Physicochemical Hydrodynamics* (Prentice-Hall, Englewood Cliffs, N.J., 1962).
- [6] V. A. Murtsovkin, *Colloid J.* **58**, 341 (1996).
- [7] N. I. Gamayunov and V. A. Murtsovkin, *Colloid J. USSR* **49**, 543 (1987); V. A. Murtsovkin and G. I. Mantrov, *Colloid J. USSR* **52**, 933 (1990).
- [8] R. Aris, *Vectors, Tensors, and the Basic Equations of Fluid Mechanics* (Dover, New York, 1962).
- [9] J. Happel and H. Brenner, *Low Reynolds Number Hydrodynamics* (Prentice-Hall, Englewood Cliffs, N.J., 1965).
- [10] H. Brenner, *Chem. Eng. Sci.* **19**, 703 (1964).
- [11] J. L. Anderson, *J. Colloid Interface Sci.* **105**, 45 (1985).